

# CONGRUENCES AND RELATIONS FOR $r$ -FISHBURN NUMBERS

F. G. GARVAN

ABSTRACT. Recently Andrews and Sellers proved some amazing congruences for the Fishburn numbers. We extend their results to a more general sequence of numbers. As a result we prove a new congruence mod 23 for the Fishburn numbers and prove their conjectured mod 5 congruence for a related sequence. We also extend and prove some unpublished conjectures of Garthwaite and Rhoades.

## 1. INTRODUCTION

The Fishburn numbers  $\xi(n)$  [2] are defined by the formal power series

$$(1.1) \quad \sum_{n=0}^{\infty} \xi(n)q^n = F(1 - q),$$

where

$$(1.2) \quad F(q) := \sum_{n=0}^{\infty} (q; q)_n,$$

and

$$(a; q)_n = (1 - a)(1 - aq) \cdots (1 - aq^{n-1}).$$

Zagier [10] showed that  $\xi(n)$  is the number of linearized chord diagrams of degree  $n$ , and also the number of nonisomorphic interval orders on  $n$  unlabeled points. Andrews and Sellers [5] proved some amazing congruences for the Fishburn numbers. For example, for all  $n \geq 0$ ,

$$(1.3) \quad \xi(5n + 3) \equiv \xi(5n + 4) \equiv 0 \pmod{5},$$

$$(1.4) \quad \xi(7n + 6) \equiv 0 \pmod{7},$$

$$(1.5) \quad \xi(11n + 8) \equiv \xi(11n + 9) \equiv \xi(11n + 10) \equiv 0 \pmod{11},$$

$$(1.6) \quad \xi(17n + 16) \equiv 0 \pmod{17}, \text{ and}$$

$$(1.7) \quad \xi(19n + 17) \equiv \xi(19n + 18) \equiv 0 \pmod{19}.$$

In fact, they prove that there are analogous congruences for all primes  $p$  that are quadratic nonresidues mod 23. For  $p$  prime they define

$$(1.8) \quad S(p) = \{j : 0 \leq j \leq p - 1 \text{ such that } \frac{1}{2}n(3n - 1) \equiv j \pmod{p} \text{ for some } n\}$$

and

$$(1.9) \quad T(p) = \{k : 0 \leq k \leq p - 1 \text{ such that } k \text{ is larger than every element of } S(p)\}.$$

---

*Date:* June 20, 2014.

*2010 Mathematics Subject Classification.* 05A19, 11B65, 11P83.

*Key words and phrases.* Fishburn numbers, interval orders, congruence, Bernoulli polynomial, Stirling numbers, Glaisher T-numbers.

We state their main result.

**Theorem 1.1** (Andrews and Sellers[5]). *If  $p$  is a prime and  $i \in T(p)$  (as defined in (1.9)), then for all  $n \geq 0$ ,*

$$\xi(pn + i) \equiv 0 \pmod{p}.$$

*Remark 1.2.* Congruences (1.3)–(1.7) are the cases  $p = 5, 7, 11, 17$  and  $19$  of Theorem 1.1. Andrews and Sellers proved that  $T(p)$  is nonempty whenever  $p$  is a quadratic nonresidue mod 23.

Recently Garthwaite and Rhoades [7] observed congruence pairs and triples such as

$$(1.10) \quad \xi(5n + 2) - 2\xi(5n + 1) \equiv 0 \pmod{5},$$

$$(1.11) \quad \xi(11n + 7) - 3\xi(11n + 4) + 2\xi(11n + 3) \equiv 0 \pmod{11}.$$

In Theorem 1.3 and Corollary 1.7 below we prove congruences relations mod  $p$  exist for all primes  $\geq 5$ .

We extend the Andrews and Sellers result to what we call  $r$ -Fishburn numbers  $\xi_r(n)$  and which we define by the formal power series

$$(1.12) \quad \sum_{n=0}^{\infty} \xi_r(n)q^n = F((1 - q)^r),$$

where  $r$  is any nonzero integer. The case  $r = 1$  corresponds to the ordinary Fishburn numbers. Even in the case  $r = 1$  we are able to augment the set  $T(p)$ . For  $p \geq 5$  prime,  $r$  relatively prime to  $p$  and nonzero and  $0 \leq s \leq p - 1$  we define

$$(1.13) \quad S^*(p, r, s) = \left\{ j : 0 \leq j \leq p - 1 \text{ such that } \frac{1}{2}rn(3n - 1) \equiv j - s \pmod{p} \text{ for some } n \right. \\ \left. \text{and } 24(j - s) \not\equiv -r \pmod{p} \right\}$$

and

$$(1.14) \quad T^*(p, r, s) = \{ k : 0 \leq k \leq p - 1 \text{ such that } k \text{ is larger than every element of } S^*(p, r, s) \}.$$

We state our main

**Theorem 1.3.** *Suppose  $p \geq 5$  is prime,  $r$  is a nonzero integer relatively prime to  $p$  and  $0 \leq s \leq p - 1$ . If  $m \in T^*(p, r, s)$  (as defined in (1.14)), then for all  $n \geq 0$ ,*

$$\sum_{j=0}^s \binom{s}{j} (-1)^j \xi_r(pn + m - j) \equiv 0 \pmod{p}.$$

*Remark 1.4.* We make some remarks.

- (i) The  $(r, s) = (1, 0)$  case of the theorem is slightly stronger than the Andrews-Sellers result. We observe that although  $22 \in S(23)$ ,  $22 \notin S^*(23, 1, 0)$  and we find  $T^*(23, 1, 0) = \{18, 19, 20, 21, 22\}$  so that

$$(1.15) \quad \xi(23n + 18) \equiv \xi(23n + 19) \equiv \xi(23n + 20) \equiv \xi(23n + 21) \equiv \xi(23n + 22) \equiv 0 \pmod{23}.$$

This is a congruence that Andrews and Sellers missed.

(ii) We find that  $S^*(5, -1, 0) = \{0, 3\}$  and  $T^*(5, -1, 0) = \{4\}$  so that

$$(1.16) \quad \xi_{-1}(5n + 4) \equiv 0 \pmod{5}.$$

This congruence was conjectured by Andrews and Sellers [5]. When  $(r, s) = (-1, 0)$  the Theorem only gives a congruence in the case  $p = 5$ . This is because 1 is a pentagonal number so that when  $p > 5$  we have  $p - 1 \in S^*(p, -1)$  and  $T^*(p, -1, 0)$  is empty. Using the facts that

$$T^*(5, -1, 2) = \{3, 4\}, \quad T^*(5, -1, 3) = \{4\},$$

we find that

$$(1.17) \quad \xi_{-1}(5n + 3) \equiv 3\xi_{-1}(5n + 2) \equiv 2\xi_{-1}(5n + 1) \pmod{5}.$$

**Example 1.5.**

$$S^*(43, -1, 2) = \{0, 1, 2, 5, 7, 10, 13, 14, 16, 18, 19, 23, 29, 30, 31, 33, 37, 38, 39, 40, 41\},$$

so that

$$T^*(43, -1, 2) = \{42\},$$

and

$$\xi_{-1}(43n + 42) - 2\xi_{-1}(43n + 41) + \xi_{-1}(43n + 40) \equiv 0 \pmod{43},$$

for all  $n \geq 0$ .

We highlight the  $s = 0$  case of the theorem.

**Corollary 1.6.** *Suppose  $p \geq 5$  is prime and  $r$  is a nonzero integer relatively prime to  $p$ . If  $m \in T^*(p, r, 0)$  (as defined in (1.14)), then for all  $n \geq 0$ ,*

$$\xi_r(pn + m) \equiv 0 \pmod{p}.$$

**Corollary 1.7.** *Suppose  $p \geq 5$  is prime, and  $r$  is a nonzero integer relatively prime to  $p$ . Then there are at least  $\frac{1}{2}(p+1)$  linearly independent congruence relations mod  $p$  of the form*

$$\sum_{j=0}^{p-1} \alpha_j \xi_r(pn + j) \equiv 0 \pmod{p},$$

where  $n$  is any nonnegative integer and  $\vec{\alpha} \in \mathbb{F}_p^p$ .

*Remark 1.8.* In Section 4 we prove Corollary 1.7 by showing that the relations

$$\sum_{j=0}^s \binom{s}{j} (-1)^j \xi_r(pn + p - 1 - j) \equiv 0 \pmod{p},$$

where the Legendre symbol  $\left(\frac{-24(1+s)\bar{r}+1}{p}\right) = -1$  or 0, form a set of  $\frac{1}{2}(p+1)$  linearly independent congruence relations mod  $p$ . Here  $r\bar{r} \equiv 1 \pmod{p}$ . This also means that  $s$  can never equal  $p - 1$ .

**Example 1.9.** When  $p = 7$  and  $r = 1$  there are 4 relations mod 7:

$$\begin{aligned}\xi(7n + 6) &\equiv 0 \pmod{7}, \\ \xi(7n + 6) - 2\xi(7n + 5) + \xi(7n + 4) &\equiv 0 \pmod{7}, \\ \xi(7n + 6) - 3\xi(7n + 5) + 3\xi(7n + 4) - \xi(7n + 3) &\equiv 0 \pmod{7}, \\ \xi(7n + 6) - 4\xi(7n + 5) + 6\xi(7n + 4) - 4\xi(7n + 3) + \xi(7n + 2) &\equiv 0 \pmod{7},\end{aligned}$$

which can be rewritten as

$$\begin{aligned}\xi(7n + 6) &\equiv 0 \pmod{7}, \\ \xi(7n + 5) + 5\xi(7n + 2) &\equiv 0 \pmod{7}, \\ \xi(7n + 4) + 3\xi(7n + 2) &\equiv 0 \pmod{7}, \\ \xi(7n + 3) + \xi(7n + 2) &\equiv 0 \pmod{7}.\end{aligned}$$

**Conjecture 1.10.** Suppose  $p \geq 5$  is prime, and  $r$  is a nonzero integer relatively prime to  $p$ . Then there are exactly  $\frac{1}{2}(p+1)$  linearly independent congruence relations mod  $p$  of the form

$$\sum_{j=0}^{p-1} \alpha_j \xi_r(pn + j) \equiv 0 \pmod{p},$$

where  $n$  is any nonnegative integer and  $\vec{\alpha} \in \mathbb{F}_p^p$ .

Following Andrews and Sellers [5], we define

$$(1.18) \quad F(q, N) = \sum_{n=0}^N (q; q)_n,$$

and the  $p$ -dissection

$$(1.19) \quad F(q, N) = \sum_{i=0}^{p-1} q^i A_p(N, i, q^p).$$

We consider the coefficients of the polynomials

$$(1.20) \quad A_p(pn - 1, i, 1 - q) = \sum_{k \geq 0} \alpha(p, n, i, k) q^k.$$

The Andrews-Sellers Theorem 1.1 depends crucially on

**Lemma 1.11** (Andrews and Sellers [5]). *If  $i \notin S(p)$ , then*

$$\alpha(p, n, i, k) = 0,$$

for  $0 \leq k \leq n - 1$ .

We consider the analog of this result when  $24i \equiv -1 \pmod{p}$ . For  $p \geq 5$  prime we define  $\bar{\xi}_p(n)$  by the formal power series

$$(1.21) \quad \sum_{n=0}^{\infty} \bar{\xi}_p(n) q^n = (1 - q)^{\lfloor \frac{p}{24} \rfloor} F((1 - q)^p),$$

where  $F(q)$  is defined in (1.2). Observe that when  $5 \leq p \leq 23$ ,

$$\bar{\xi}_p(n) = \xi_p(n),$$

for  $n \geq 0$ . We find the following new relation for Fishburn numbers.

**Theorem 1.12.** *Suppose  $p \geq 5$  is prime and  $24i_0 \equiv -1 \pmod{p}$  where  $1 \leq i_0 \leq p-1$ . Then*

$$\alpha(p, n, i_0, k) = p \binom{12}{p} \bar{\xi}_p(k),$$

for  $0 \leq k \leq n-1$ . Here  $\binom{\cdot}{\cdot}$  is the Kronecker symbol.

Our main Theorem 1.3 will follow from Lemma 1.11 and Theorem 1.12 in a straightforward manner.

## 2. PRELIMINARY RESULTS

**Lemma 2.1.** *Let  $p$  be prime and suppose  $0 \leq j \leq p-1$  and  $0 \leq k \leq M-1 \leq N-1$ . Then*

$$\alpha(p, N, j, k) = \alpha(p, M, j, k),$$

where  $\alpha(p, n, i, k)$  is defined in (1.20).

*Proof.* Let  $\zeta = \exp(2\pi i/p)$ . Then from (1.19) we have

$$A_p(N, j, q) = \frac{1}{p} \sum_{k=0}^{p-1} \zeta^{-jk} q^{-\frac{j}{p}} F\left(\zeta^k q^{\frac{1}{p}}, N\right).$$

Next we suppose that  $n \geq pM$ . Then

$$\begin{aligned} \left(\zeta^k(1-q)^{\frac{1}{p}}; \zeta^k(1-q)^{\frac{1}{p}}\right)_n &= \prod_{j=1}^n \left(1 - \left(\zeta^k(1-q)^{\frac{1}{p}}\right)^j\right) \\ &= \prod_{j=1}^{\lfloor n/p \rfloor} (1 - (1-q)^j) \prod_{\substack{j=1 \\ j \not\equiv 0 \pmod{p}}}^n \left(1 - \left(\zeta^k(1-q)^{\frac{1}{p}}\right)^j\right) \\ &= \prod_{j=1}^M (jq + O(q^2)) \prod_{\substack{j=1 \\ j \not\equiv 0 \pmod{p}}}^n \left(1 - \left(\zeta^k(1-q)^{\frac{1}{p}}\right)^j\right) \\ &= O(q^M). \end{aligned}$$

Thus

$$\begin{aligned} A_p(pN-1, j, 1-q) &= \frac{1}{p} \sum_{k=0}^{p-1} \zeta^{-jk} q^{-\frac{j}{p}} \sum_{n=0}^{pN-1} \left(\zeta^k(1-q)^{\frac{1}{p}}; \zeta^k(1-q)^{\frac{1}{p}}\right)_n \\ &= A_p(pM-1, j, 1-q) + O(q^M). \end{aligned}$$

The result follows.  $\square$

Andrews and Sellers define a Stirling like array of numbers  $C(n, i, j, p)$  for  $n \geq 0$ ,  $0 \leq i \leq p - 1$ , and  $0 \leq j \leq n$ , which are defined by the recursion

$$(2.1) \quad C(n+1, i, j, p) = (i + jp)C(n, i, j, p) + pC(n, i, j-1, p),$$

and the initial value

$$(2.2) \quad C(0, i, 0, p) = 1.$$

It is understood that if either of the conditions  $n \geq 0$  or  $0 \leq j \leq n$  are not satisfied, then  $C(n, i, j, p) = 0$ . We note that

$$C(n, i, 0, p) = i^n.$$

We need a generalization of the signless Stirling numbers of the first kind. We define the numbers  $s_1(n, j, m)$  for  $0 \leq j \leq n$  by

$$(2.3) \quad \sum_{j=0}^n s_1(n, j, m)x^j = (x-m)(x-m+1) \cdots (x-m+n-1).$$

We note case  $m = 0$  correspond to the signless Stirling numbers of the first kind  $s_1(n, j)$ . We define

$$(2.4) \quad f(x, n, k, m) = (-1)^n \sum_{j=k}^n \binom{j}{k} s_1(n, j, m)x^j,$$

for  $0 \leq k \leq n$ , otherwise define  $f(x, n, k, m) = 0$ .

**Lemma 2.2.** For  $0 \leq k \leq n+1$  we have

$$(2.5) \quad f(x, n+1, k, m) = -((x+n-m)f(x, n, k, m) + xf(x, n, k-1, m)).$$

*Proof.* First we observe that

$$f(x, n, k, m) = \frac{x^k}{k!} \left( \frac{\partial}{\partial x} \right)^k f(x, n, 0, m),$$

where

$$f(x, n, 0, m) = (-1)^n (x-m)(x-m+1) \cdots (x-m+n-1) \quad (\text{by (2.3)}),$$

for  $k \geq 0$ . Now we let

$$\tilde{f}(x, n, k, m) = \left( \frac{\partial}{\partial x} \right)^k (-1)^n (x-m)(x-m+1) \cdots (x-m+n-1),$$

for  $0 \leq k \leq n$ , otherwise define  $\tilde{f}(x, n, k, m) = 0$ . We show that

$$(2.6) \quad \tilde{f}(x, n+1, k, m) = -\left( (x-m+n)\tilde{f}(x, n, k, m) + k\tilde{f}(x, n, k-1, m) \right)$$

where  $0 \leq k \leq n+1$ . Since

$$\tilde{f}(x, n+1, 0, m) = -(x-m+n)\tilde{f}(x, n, 0, m),$$

we have

$$\frac{\partial}{\partial x} \tilde{f}(x, n+1, 0, m) = -(x-m+n) \frac{\partial}{\partial x} \tilde{f}(x, n, 0, m) - \tilde{f}(x, n, 0, m),$$

so that

$$\tilde{f}(x, n + 1, 1, m) = - \left( (x - m + n) \tilde{f}(x, n, 1, m) + \tilde{f}(x, n, 0, m) \right)$$

and (2.6) holds for  $k = 1$ . We assume (2.6) holds for  $k \leq K$ .

$$\begin{aligned} \tilde{f}(x, n + 1, K + 1, m) &= \frac{\partial}{\partial x} \tilde{f}(x, n + 1, K, m) \\ &= - \frac{\partial}{\partial x} \left[ (x - m + n) \tilde{f}(x, n, K, m) + K \tilde{f}(x, n, k - 1, m) \right] \\ &= - \left[ (x - m + n) \tilde{f}(x, n, K + 1, m) + \tilde{f}(x, n, K, m) + K \tilde{f}(x, n, K, m) \right] \\ &= - \left[ (x - m + n) \tilde{f}(x, n, K + 1, m) + (K + 1) \tilde{f}(x, n, K, m) \right], \end{aligned}$$

and (2.6) holds for  $k = K + 1$ . Hence (2.6) holds for all  $k$  by induction. Since

$$f(x, n, k, m) = \frac{x^k}{k!} \tilde{f}(x, n, k, m),$$

the result (2.5) follows easily. □

**Theorem 2.3.** Let  $i_0 = (p^2 - 1)z - mp$ . Suppose that

$$(2.7) \quad \sum_{\ell=0}^n C(n, i_0, \ell, p) A_1(\ell, m) = (-1)^n z^n \sum_{k=0}^n \binom{n}{k} X(k),$$

for  $n \geq 0$ . Then

$$(2.8) \quad A_1(n, m) = (-1)^n \sum_{k=0}^n \sum_{j=k}^n \binom{j}{k} s_1(n, j, m) p^{j-2k} X(k) z^j,$$

for  $n \geq 0$ .

*Proof.* Since for  $n \geq 0$  (2.7) forms a triangular system of equations in the unknowns  $A_1(\ell, m)$  (for fixed  $m$ ) and each diagonal coefficient

$$C(n, i_0, n, p) = p^n \neq 0,$$

it suffices to show that  $A_1(n, m)$  given by (2.8) satisfies (2.7). By considering the coefficient of  $X(L)$  it suffices to show that

$$(2.9) \quad \sum_{\ell=0}^n C(n, i_0, \ell, p) f(pz, \ell, L, m) = G(n, L),$$

where

$$G(n, L) = (-1)^n \binom{n}{L} z^n p^{2L},$$

for  $0 \leq L \leq n$  and  $m \geq 0$ . We proceed by induction on  $n$ . The result is clearly true for  $n = 0$ . We assume (2.9) holds for  $n = N$ . Now

$$\sum_{\ell=0}^{N+1} C(N + 1, i_0, \ell, p) f(pz, \ell, L, m)$$

$$\begin{aligned}
&= \sum_{\ell=0}^{N+1} (((p^2 - 1)z - mp + \ell p)C(N, i_0, \ell, p) + pC(N, i_0, \ell - 1, p)) f(pz, \ell, L, m) \quad (\text{by (2.1)}) \\
&= \sum_{\ell=0}^N ((p^2 - 1)z - mp + \ell p)C(N, i_0, \ell, p) f(pz, \ell, L, m) \\
&\quad + \sum_{\ell=0}^N pC(N, i_0, \ell, p) f(pz, \ell + 1, L, m) \\
&= \sum_{\ell=0}^N ((p^2 - 1)z - mp + \ell p)C(N, i_0, \ell, p) f(pz, \ell, L, m) \\
&\quad - \sum_{\ell=0}^N pC(N, i_0, \ell, p) \left( (pz + \ell - m) f(pz, \ell, L, m) + pz f(pz, \ell, L - 1, m) \right) \\
&= -z \sum_{\ell=0}^N C(N, i_0, \ell, p) f(pz, \ell, L, m) - p^2 z \sum_{\ell=0}^N C(N, i_0, \ell, p) f(pz, \ell, L - 1, m) \\
&\qquad\qquad\qquad (\text{by (2.5)}) \\
&= -zG(N, L) - p^2 zG(N, L - 1) \\
&= (-1)^{N+1} \left( \binom{N}{L} z^{N+1} p^{2L} + \binom{N}{L-1} z^{N+1} p^{2L} \right) \\
&= (-1)^{N+1} \binom{N+1}{L} z^{N+1} p^{2L} = G(N+1, L),
\end{aligned}$$

and (2.9) holds for  $n = N + 1$ , thus completing our induction proof.  $\square$

We will also need some results of Zagier [10] on the Fishburn numbers. We need the following formal power series identity [10, Eqn.(4),p.946]

$$(2.10) \quad e^{t/24} \sum_{n=0}^{\infty} (1 - e^t) \cdots (1 - e^{nt}) = \sum_{n=0}^{\infty} \frac{T_n}{n!} \left( \frac{-t}{24} \right)^n,$$

where  $T_n$  are the Glaisher  $T$ -numbers [1] and which are given explicitly by

$$(2.11) \quad T_n = 6 \frac{(-144)^n}{n+1} \left[ B_{2n+2} \left( \frac{1}{12} \right) - B_{2n+2} \left( \frac{5}{12} \right) \right],$$

where  $B_n(x)$  denotes the  $n$ -th Bernoulli polynomial. We remark that letting  $t = \log(1 - q)$  in (2.10) and using (3.10) below we find that

$$(2.12) \quad \xi(n) = \sum_{m=0}^n \sum_{k=0}^m (-1)^{n+k} \binom{-1/24}{n-m} \frac{s_1(m, k)}{m! 24^k} T_k,$$



which is useful for calculation. Zagier [10] also determined the behaviour of  $F(q)$  when  $q$  is near a root of unity. In particular, if  $\zeta = \zeta_p$  is a  $p$ -th root of unity and  $N = 12p$  then

$$(2.13) \quad e^{t/24} F(\zeta e^t) = \sum_{n=0}^{\infty} \frac{c_n(\zeta)}{n!} \left( \frac{-t}{24} \right)^n,$$

where

$$(2.14) \quad c_n(\zeta) = \frac{(-1)^n N^{2n+1}}{2n+2} \sum_{m=1}^{N/2} \chi(m) \zeta^{\frac{1}{24}(m^2-1)} B_{2n+2} \left( \frac{m}{N} \right),$$

and where  $\chi$  is the character mod 12 given by

$$(2.15) \quad \chi(n) = \left( \frac{12}{n} \right) = \begin{cases} 1 & \text{if } n \equiv \pm 1 \pmod{12}, \\ -1 & \text{if } n \equiv \pm 5 \pmod{12}, \\ 0 & \text{otherwise.} \end{cases}$$

This character occurs in the statement of Theorem 1.12.

### 3. PROOF OF THEOREM 1.12

In this section we assume  $p > 3$  is prime,  $N = 12p$  and  $\zeta$  is any  $p$ -th root of unity. Following [10] we define the sequence  $(b_n(\zeta))$  formally by

$$(3.1) \quad F(\zeta e^t) = \sum_{n=0}^{\infty} \frac{b_n(\zeta)}{n!} t^n.$$

From (2.10), (2.13), (2.14) we have for  $n \geq 0$

$$(3.2) \quad b_n(\zeta) = \frac{(-1)^n}{24^n} \sum_{j=0}^n \binom{n}{j} \sum_{i=0}^{p-1} \gamma(j, i) \zeta^i,$$

where

$$(3.3) \quad \gamma(j, i) = \frac{(-1)^j N^{2j+1}}{2j+2} \sum_{\substack{m=1 \\ (m^2-1)/24 \equiv i \pmod{p}}}^{N/2} \chi(m) B_{2j+2} \left( \frac{m}{N} \right).$$

As in [5] we have for  $n \geq 0$

$$b_n(\zeta) = \left( \frac{d}{dt} \right)^n F(\zeta e^t) \Big|_{t=0} = \left( q \frac{d}{dq} \right)^n F(q) \Big|_{q=\zeta} = \left( q \frac{d}{dq} \right)^n F(q, m) \Big|_{q=\zeta}$$

for  $m \geq (n+1)p - 1$ . Proceeding as in the proof of [5, lemma 2.5] we have from (2.14), (3.2) and Lemma 2.1 that

$$\begin{aligned} b_n(\zeta) &= \left( q \frac{d}{dq} \right)^n F(q, (n+1)p - 1) \Big|_{q=\zeta} \\ &= \sum_{j=0}^n \sum_{i=0}^{p-1} C(n, i, j, p) \zeta^i A_p^{(j)}(p(n+1) - 1, i, 1) \end{aligned}$$

$$\begin{aligned}
&= \sum_{j=0}^n \sum_{i=0}^{p-1} C(n, i, j, p) \zeta^i A_p^{(j)}(p(j+1) - 1, i, 1) \\
&= \frac{(-1)^n}{24^n} \sum_{j=0}^n \binom{n}{j} \sum_{i=0}^{p-1} \gamma(j, i) \zeta^i.
\end{aligned}$$

Since this identity holds for all  $p$ -th roots of unity  $\zeta$  (including  $\zeta = 1$ ) and all the coefficients involved are all rational numbers we may equate coefficients of  $\zeta^i$  on both sides to obtain

$$(3.4) \quad \sum_{j=0}^n C(n, i, j, p) A_p^{(j)}(p(j+1) - 1, i, 1) = \frac{(-1)^n}{24^n} \sum_{j=0}^n \binom{n}{j} \gamma(j, i),$$

for  $0 \leq i \leq p-1$ . Now we let  $i_0$  be the least nonnegative integer satisfying  $24i_0 \equiv -1 \pmod{p}$ . We find that

$$(3.5) \quad i_0 = \frac{p^2 - 1}{24} - \left\lfloor \frac{p}{24} \right\rfloor p.$$

We now calculate  $\gamma(j, i_0)$ . We see that

$$\frac{m^2 - 1}{24} \equiv i_0 \pmod{p} \text{ if and only if } m \equiv 0 \pmod{p}.$$

In the sum (3.3) (with  $(N = 12p)$ ) we only consider the terms with  $m = p, 5p$  to find that

$$(3.6) \quad \gamma(j, i_0) = \chi(p) \frac{(-1)^j 12^{2j+1} p^{2j+1}}{2j+2} \left( B_{2j+2} \left( \frac{1}{12} \right) - B_{2j+2} \left( \frac{5}{12} \right) \right).$$

We apply Theorem 2.3 with  $z = \frac{1}{24}$  and  $m = \left\lfloor \frac{p}{24} \right\rfloor$  to equation (3.4) with  $i = i_0$  to obtain

$$\begin{aligned}
A_p^{(n)}(p(n+1) - 1, i_0, 1) &= (-1)^n \sum_{k=0}^n \sum_{j=k}^n \binom{j}{k} s_1(n, j, \lfloor p/24 \rfloor) \frac{p^{j+1}}{24^j} \\
&\quad \chi(p) \frac{(-1)^k 12^{2k+1}}{2k+2} \left( B_{2k+2} \left( \frac{1}{12} \right) - B_{2k+2} \left( \frac{5}{12} \right) \right) \\
&= (-1)^n \chi(p) \sum_{k=0}^n \sum_{j=k}^n \binom{j}{k} s_1(n, j, \lfloor p/24 \rfloor) \frac{p^{j+1}}{24^j} T_k,
\end{aligned}$$

by (2.11). From (1.20) we see that

$$\alpha(p, n+1, i_0, n) = \frac{(-1)^n}{n!} A_p^{(n)}(p(n+1) - 1, i_0, 1).$$

Hence

$$\begin{aligned}
\sum_{n=0}^{\infty} \alpha(p, n+1, i_0, n) x^n &= p \chi(p) \sum_{n=0}^{\infty} \frac{x^n}{n!} \sum_{k=0}^n \sum_{j=k}^n \binom{j}{k} s_1(n, j, \lfloor p/24 \rfloor) \frac{p^j}{24^j} T_k, \\
(3.7) \quad &= p \chi(p) \sum_{j=0}^{\infty} \sum_{k=0}^j \left( \sum_{n=j}^{\infty} s_1(n, j, \lfloor p/24 \rfloor) \frac{x^n}{n!} \right) \frac{p^j}{24^j} \binom{j}{k} T_k
\end{aligned}$$

It is well-known that the signless Stirling numbers of the first kind have generating function

$$(3.8) \quad G(x, u) = \exp(-u \log 1 - x) = (1 - x)^{-u} = \sum_{n=0}^{\infty} \sum_{j=0}^n s_1(n, j) u^j \frac{x^n}{n!}.$$

Our generalized signless Stirling numbers have generating function

$$(3.9) \quad G_k(x, u) = (1 - x)^k \exp(-u \log 1 - x) = (1 - x)^{k-u} = \sum_{n=0}^{\infty} \sum_{j=0}^n s_1(n, j, k) u^j \frac{x^n}{n!}.$$

Thus

$$(3.10) \quad \sum_{n=j}^{\infty} s_1(n, j, k) \frac{x^n}{n!} = (1 - x)^k \frac{(-\log(1 - x))^j}{j!}.$$

Hence from (3.10) and (3.7) we have

$$(3.11) \quad \sum_{n=0}^{\infty} \alpha(p, n + 1, i_0, n) q^n = p\chi(p)(1 - q)^{\lfloor p/24 \rfloor} \sum_{j=0}^{\infty} \left( \sum_{k=0}^j \binom{j}{k} T_k \right) \frac{(-p \log(1 - q))^j}{24^j j!}$$

From Zagier's result (2.10) we have

$$(3.12) \quad F(\exp(t)) = \sum_{n=0}^{\infty} \left( \sum_{k=0}^n \binom{n}{k} T_k \right) \frac{(-t)^n}{24^n n!}.$$

Thus using this result in (3.11) we have

$$(3.13) \quad \begin{aligned} \sum_{n=0}^{\infty} \alpha(p, n + 1, i_0, n) q^n &= p\chi(p)(1 - q)^{\lfloor p/24 \rfloor} F(\exp(p \log(1 - q))) \\ &= p\chi(p)(1 - q)^{\lfloor p/24 \rfloor} F((1 - q)^p). \end{aligned}$$

Theorem 1.12 follows from (1.21) and (3.13) since

$$\alpha(p, n, i_0, k) = \alpha(p, k + 1, i_0, k),$$

for  $0 \leq k \leq n - 1$  by Lemma 2.1.

#### 4. PROOF OF THEOREM 1.3 AND COROLLARY 1.7

We assume  $p \geq 5$  is prime and  $24i_0 \equiv -1 \pmod{p}$  where  $1 \leq i_0 \leq p - 1$ . For two formal power series

$$A = \sum_{n=0}^{\infty} a_n q^n, \quad B = \sum_{n=0}^{\infty} b_n q^n \in \mathbb{Z}[[q]],$$

we write

$$A \equiv B \pmod{p} \quad \text{if} \quad a_n \equiv b_n \pmod{p},$$

for all  $n \geq 0$ . If  $p$  is prime and  $m$  is a nonnegative integer

$$m = jp + r, \quad \text{where } 0 \leq r < p \text{ and } j, r \in \mathbb{N},$$

then

$$(4.1) \quad (1 - q)^m = (1 - q)^r (1 - q)^{pj} \equiv (1 - q)^r (1 - q^p)^j \pmod{p}.$$

Now suppose  $r$  is a nonzero integer relatively prime to  $p$  and  $0 \leq s \leq p-1$ . We consider two cases.

**Case I.**  $r > 0$ . We proceed as in [5, Section 3]. We note that

$$(4.2) \quad \sum_{n=0}^{\infty} \xi(n)q^n = F(1-q, N) + O(q^{N+1}).$$

Now from (1.19) we have

$$\begin{aligned} F(q, pn-1) &= \sum_{i=0}^{p-1} q^i A_p(pn-1, i, q^p) \\ &= \sum_{i \in S(p) \setminus \{i_0\}} q^i A_p(pn-1, i, q^p) + q^{i_0} A_p(pn-1, i_0, q^p) \\ &\quad + \sum_{i \notin S(p)} q^i A_p(pn-1, i, q^p). \end{aligned}$$

Hence

$$\begin{aligned} F((1-q)^r, pn-1) &= \sum_{i \in S(p) \setminus \{i_0\}} (1-q)^{ri} A_p(pn-1, i, (1-q)^{rp}) \\ &\quad + (1-q)^{ri_0} A_p(pn-1, i_0, (1-q)^{rp}) \\ &\quad + \sum_{i \notin S(p)} (1-q)^{ri} A_p(pn-1, i, (1-q)^{rp}). \end{aligned}$$

For  $i \notin S(p)$  we have

$$A_p(pn-1, i, q) = \sum_{k \geq 0} \alpha(p, n, i, k)(1-q)^k,$$

and

$$\begin{aligned} A_p(pn-1, i, (1-q)^{rp}) &= \sum_{k \geq 0} \alpha(p, n, i, k)(1 - (1-q)^{rp})^k \\ &\equiv \sum_{k \geq 0} \alpha(p, n, i, k)(1 - (1-q^p)^r)^k \pmod{p} \\ &\equiv O(q^{pn}) \pmod{p}, \end{aligned}$$

by Lemma 1.11. In a similar fashion we have

$$A_p(pn-1, i_0, (1-q)^{rp}) \equiv O(q^{pn}) \pmod{p},$$

using Theorem 1.12. Thus

$$(1-q)^s F((1-q)^r, pn-1) \equiv \sum_{i \in S(p) \setminus \{i_0\}} (1-q)^{ri+s} A_p(pn-1, i, (1-q^p)^r) + O(q^{pn}) \pmod{p}.$$

By (4.1) we see that the only terms  $q^{j'}$  that occur in  $(1-q)^{ri+s}$  (where  $i \in S(p) \setminus \{i_0\}$ ) satisfy

$$j' \equiv j \pmod{p} \quad \text{where } 0 \leq j \leq m \text{ and } m \in S^*(p, r, s).$$

This is because  $i \in S(p) \setminus \{i_0\}$  if and only if  $ri + s$  is congruent to an element of  $S^*(p, r, s)$ . Since  $A_p(pn - 1, i, (1 - q^p)^r)$  is a polynomial in  $q^p$  the result follows by letting  $n \rightarrow \infty$ ; i.e. every term  $q^{j'}$  in

$$(1 - q)^s F((1 - q)^r) = \sum_{n=0}^{\infty} \sum_{j=0}^s \binom{s}{j} (-1)^j \xi_r(n - j) q^n$$

where  $j'$  is congruent to an element of  $T^*(p, r, s)$  must have a coefficient that is congruent to 0 mod  $p$ .

**Case II.**  $r < 0$ . This time we choose integers  $\beta$  and  $m$  such that

$$r = mp + \beta,$$

where  $0 < \beta \leq p - 1$  and  $m < 0$ . We find that

$$(1 - q)^r = (1 - q)^\beta \Phi(q^p) + p\Psi(q) \equiv (1 - q)^\beta \Phi(q^p) \pmod{p},$$

where  $\Phi(q), \Psi(q) \in \mathbb{Z}[[q]]$  and  $\Phi(q)$  has constant term 1 so that it is a unit in the ring of formal power series. In fact,

$$\Phi(q) = (1 - q)^m = 1 + \sum_{k=1}^{\infty} \binom{k - m - 1}{k} q^k = 1 - mq + \frac{m(m - 1)}{2} q^2 + \dots$$

We basically proceed as in Case I. We have

$$\begin{aligned} F((1 - q)^r, pn - 1) &= F((1 - q)^\beta \Phi(q^p) + p\Psi(q), pn - 1) \\ &= \sum_{i=0}^{p-1} ((1 - q)^\beta \Phi(q^p) + p\Psi(q))^i A_p(pn - 1, i, ((1 - q)^\beta \Phi(q^p) + p\Psi(q))^p) \\ &\equiv \sum_{i=0}^{p-1} (1 - q)^{\beta i} [\Phi(q^p)]^i A_p(pn - 1, i, ((1 - q^p)^\beta [\Phi(q^p)]^p) \pmod{p} \\ &\equiv \sum_{i \in S(p) \setminus \{i_0\}} (1 - q)^{\beta i} [\Phi(q^p)]^p A_p(pn - 1, i, ((1 - q^p)^\beta [\Phi(q^p)]^p) + O(q^{pn}) \pmod{p}, \end{aligned}$$

and

$$\begin{aligned} (1 - q)^s F((1 - q)^r, pn - 1) \\ \equiv \sum_{i \in S(p) \setminus \{i_0\}} (1 - q)^{\beta i + s} [\Phi(q^p)]^p A_p(pn - 1, i, ((1 - q^p)^\beta [\Phi(q^p)]^p) + O(q^{pn}) \pmod{p}, \end{aligned}$$

arguing as before. This time instead of the term

$$A_p(pn - 1, i, (1 - q^p)^r),$$

which is a polynomial in  $q^p$  with integer coefficients we have the term

$$[\Phi(q^p)]^p A_p(pn - 1, i, ((1 - q^p)^\beta [\Phi(q^p)]^p),$$

which is a formal power series in  $q^p$  with integer coefficients. The result follows as before by letting  $n \rightarrow \infty$ . This completes the proof of our main theorem.

We now prove Corollary 1.7. As before suppose  $p \geq 5$  is prime and let  $r$  be a fixed nonzero integer relatively prime to  $p$ . Suppose  $\bar{r}$  is the multiplicative inverse of  $r \pmod{p}$ . We see that

$$r^{\frac{1}{2}}n(3n-1) \equiv -1-s \pmod{p} \quad \text{if and only if} \quad (6n-1)^2 \equiv -24(1+s)\bar{r}+1 \pmod{p}.$$

Thus  $p-1 \notin S^*(p, r, s)$  if  $-24(1+s)\bar{r}+1$  is either a quadratic nonresidue mod  $p$  or congruent to zero mod  $p$ . There are  $\frac{1}{2}(p+1)$  such values of  $s$  for  $0 \leq s \leq p-1$ . Thus

$$(4.3) \quad \sum_{j=0}^s \binom{s}{j} (-1)^j \xi_r(pn+p-1-j) \equiv 0 \pmod{p},$$

for all  $n \geq 0$  provided  $\left(\frac{-24(1+s)\bar{r}+1}{p}\right) = -1$  or  $0$ . It is clear that this set of  $\frac{1}{2}(p+1)$  congruence relations mod  $p$  is linearly independent. This completes the proof of Corollary 1.7.

We illustrate (4.3) with some examples.

**Example 4.1.**  $s = 0$  We have

$$\xi(pn+p-1) \equiv 0 \pmod{p}$$

for all  $n \geq 0$  provided  $p \geq 5$  is prime and  $\left(\frac{-23}{p}\right) = 0$  or  $-1$ ; i.e.  $p \equiv 0, 5, 7, 10, 11, 14, 15, 17, 19, 20, 21, \text{ or } 22 \pmod{23}$ .

**Example 4.2.**  $s = 1$  We have

$$\xi(pn+p-1) \equiv \xi(pn+p-2) \pmod{p}$$

for all  $n \geq 0$  provided  $p \geq 5$  is prime and  $\left(\frac{-47}{p}\right) = 0$  or  $-1$ ; i.e.  $p \equiv 0, 5, 10, 11, 13, 15, 19, 20, 22, 23, 26, 29, 30, 31, 33, 35, 38, 39, 40, 41, 43, 44, 45, \text{ or } 46 \pmod{47}$ .

**Example 4.3.**  $s = 2$  We have

$$\xi(pn+p-1) - 2\xi(pn+p-2) + \xi(pn+p-3) \equiv 0 \pmod{p}$$

for all  $n \geq 0$  provided  $p \geq 5$  is prime and  $\left(\frac{-71}{p}\right) = 0$  or  $-1$ ; i.e.  $p = 71$  or  $\left(\frac{71}{p}\right) = -1$ . We have

$$\xi_{-1}(pn+p-1) - 2\xi_{-1}(pn+p-2) + \xi_{-1}(pn+p-3) \equiv 0 \pmod{p}$$

for all  $n \geq 0$  provided  $p \geq 5$  is prime and  $\left(\frac{73}{p}\right) = 0$  or  $-1$ ; i.e.  $p = 73$  or  $\left(\frac{73}{p}\right) = -1$ .

## 5. CONCLUSION

We pose the following problems.

- (i) The numbers  $\xi(n)$  and  $\xi_{-1}(n)$  have many combinatorial interpretations [6], [8], [9]. Use one of the interpretations to find a rank or crank-type function [3] to explain combinatorially the simplest congruences

$$\begin{aligned} \xi(5n+3) &\equiv \xi(5n+4) \equiv 0 \pmod{5}, \\ \xi_{-1}(5n+4) &\equiv 0 \pmod{5}. \end{aligned}$$

- (ii) Zagier [10] showed that  $q^{1/24}F(q)$  is a so-called quantum modular form [11]. Find and prove congruences for the coefficients of other quantum modular forms. In particular look at the functions considered by Andrews, Jiménez-Urroz and Ono [4].

### Acknowledgements

I would like to thank George Andrews, Sharon Garthwaite, Ira Gessel, Ken Ono and Rob Rhoades for their comments and suggestions. In particular I thank Ira Gessel for reminding me of the exponential generating for the signless Stirling numbers of the first kind, and I thank Sharon Garthwaite and Rob Rhoades for sharing their observations on congruences for pairs and triples for the sequences  $\xi(n)$  and  $\xi_{-1}(n)$ .

### REFERENCES

1. *Sequence A002439 (Glaisher's T numbers)*, in “The On-Line Encyclopedia of Integer Sequences”, published electronically at <http://oeis.org/A002439>, 2010.
2. *Sequence A022493 (The Fishburn numbers)*, in “The On-Line Encyclopedia of Integer Sequences”, published electronically at <http://oeis.org/A022493>, 2010.
3. G.E. Andrews and F. G. Garvan, *Dyson's crank of a partition*, Bull. Amer. Math. Soc. (N.S.) **18** (1988), 167–171.
4. G. E. Andrews, J. Jiménez-Urroz, and K. Ono, *q-series identities and values of certain L-functions*, Duke Math. J. **108** (2001), 395–419.
5. G. E. Andrews and J. A. Sellers, *Congruences for the Fishburn numbers*, preprint (arXiv:1401.5345).
6. K. Bringmann, Y. Li and R. Rhoades, *Asymptotics for the number of row Fishburn matrices*, J. Combin. Theory Ser. A, to appear.
7. S. A. Garthwaite and R. C. Rhoades, Private communication, March 4, 2014.
8. S. M. Khamis, *Exact counting of unlabeled rigid interval posets regarding or disregarding height*, Order **29** (2012), 443–461.
9. P. Levande, *Fishburn diagrams, Fishburn numbers and their refined generating functions*, J. Combin. Theory Ser. A **120** (2013), 194–217.
10. D. Zagier, *Vassiliev invariants and a strange identity related to the Dedekind eta-function*, Topology **40** (2001), 945–960.
11. D. Zagier, *Quantum modular forms*, in “Quanta of Maths,” Clay Math. Proc. **11**, Amer. Math. Soc., Providence, RI, (2010) 659–675.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, GAINESVILLE, FL 32611-8105  
*E-mail address:* fgarvan@ufl.edu