

```
[> with(qseries) :
=> with(ETA) :
```

```
[> gpA1 := [1, 2, 4, 3, 2, -5, 50, 5, 25, -2, 100, -3] :
=> epA1 := gp2etaprod(gpA1) ;
```

$$epA1 := \frac{\eta(\tau)^2 \eta(4\tau)^3 \eta(50\tau)^5}{\eta(2\tau)^5 \eta(25\tau)^2 \eta(100\tau)^3} \quad (1)$$

```
[> A1q := etaprodtoqseries(epA1, 1000) :
```

```
[> gpa1 := [1, 2, 10, 4, 2, -4, 5, -2] :
=> gpa1 := [1, 2, 10, 4, 2, -4, 5, -2]
=> epa1 := gp2etaprod(gpa1) ;
```

$$epa1 := \frac{\eta(\tau)^2 \eta(10\tau)^4}{\eta(2\tau)^4 \eta(5\tau)^2} \quad (2)$$

```
[> aq1 := etaprodtoqseries(epa1, 2000) :
=> gprho := [2, 2, 20, 4, 4, -4, 10, -2] :
=> eprho := gp2etaprod(gprho) ;
```

$$eprho := \frac{\eta(2\tau)^2 \eta(20\tau)^4}{\eta(4\tau)^4 \eta(10\tau)^2} \quad (3)$$

```
[> Gla := etaprodtoqseries(eprho, 1001) :
=> G1 := convert(series(1-Gla, q, 1020), polynom) :
=> etamake(G1, q, 100) ;
```

$$\frac{\eta(10\tau)^3 \eta(2\tau)}{\eta(20\tau) \eta(4\tau)^3} \quad (4)$$

```
[> U01 := sift(A1q, q, 5, 0, 2000) :
```

```
[> etacombo := findlincombo(U01, [seq(aq1^k, k=-2..5), seq(aq1^k*(G1),
k=-2..5)], [seq(etamake((aq1)^k, q, 100), k=-2..5), seq(etamake(
(aq1)^k*(G1), q, 100), k=-2..5)], q, 0) ;
```

nx = , 16

of terms , 37

$$etacombo := -\frac{53 \eta(\tau)^2 \eta(10\tau)^4}{\eta(2\tau)^4 \eta(5\tau)^2} + \frac{350 \eta(10\tau)^8 \eta(\tau)^4}{\eta(5\tau)^4 \eta(2\tau)^8} - \frac{1050 \eta(10\tau)^{12} \eta(\tau)^6}{\eta(5\tau)^6 \eta(2\tau)^{12}} \quad (5)$$

$$+ \frac{1375 \eta(10\tau)^{16} \eta(\tau)^8}{\eta(5\tau)^8 \eta(2\tau)^{16}} - \frac{625 \eta(10\tau)^{20} \eta(\tau)^{10}}{\eta(5\tau)^{10} \eta(2\tau)^{20}} + \frac{13 \eta(10\tau)^3 \eta(2\tau)}{\eta(20\tau) \eta(4\tau)^3}$$

$$- \frac{75 \eta(10\tau)^7 \eta(\tau)^2}{\eta(20\tau) \eta(5\tau)^2 \eta(4\tau)^3 \eta(2\tau)^3} + \frac{175 \eta(10\tau)^{11} \eta(\tau)^4}{\eta(20\tau) \eta(5\tau)^4 \eta(4\tau)^3 \eta(2\tau)^7}$$

$$-\frac{125 \eta(10 \tau)^{15} \eta(\tau)^6}{\eta(20 \tau) \eta(5 \tau)^6 \eta(4 \tau)^3 \eta(2 \tau)^{11}}$$

> #read pf:

> provemodfuncGAMMA0UpETAid(epA1,5,etacombo,20);

*** There were NO errors.

*** o EP is an MF on Gamma[0](100)

*** o Each term in the etacombo is a modular function on Gamma0(20).

*** o We also checked that the total order of each term etacombo was zero.

*** To prove the identity $U[5](EP)=etacombo$ we need to show that $v[oo](ID) > 9$ This means checking up to $q^{(10)}$.

Do you want to prove the identity? (yes/no)

You entered yes.

We verify the identity to $O(q^{(49)})$.

We find that LHS - RHS is

$$O(q^{49})$$

RESULT: The identity holds to $O(q^{(49)})$.

CONCLUSION: This proves the identity since we had only to show that $v[oo](ID) > 9$.